


Space weather and solar-terrestrial relations



solar eclipse, 11.8.1999, Wendy Carlos and John Kern



MAX-PLANCK-GESELLSCHAFT

Hardi Peter



Early note on solar-terrestrial relations

from *Richard A. Proctor:*

“Other Worlds Than Ours”, 1870. Chapter II. What we Learn From the Sun.

[In] 1859, the eminent solar observer, Carrington noticed the apparition of a **bright spot upon the Sun's surface**.


The light of this spot was so intense that he imagined the screen which shaded the plate employed to receive the solar image had been broken. (...)

Now it was found that the self-registering **magnetic** instruments of the Kew observatory had been sharply disturbed at the instant when the bright spot was seen. (...)

Telegraphic communication was interrupted, and at a station in Norway the telegraphic **apparatus was set on fire**;

auroras appeared both in the northern and southern hemispheres during the night which followed.

What is space weather?



Space weather happens when a solar storm from the Sun travels through space and impacts the Earth's magnetosphere.


Studying space weather is important to our national economy because solar storms can affect the advanced technology we have become so dependent upon in our everyday lives.

Energy and radiation from solar flares and coronal mass ejections can


- Harm astronauts in space
- Damage sensitive electronics on orbiting spacecraft...
- Cause colorful auroras, often seen in the higher latitudes...
- Create blackouts on Earth when they cause surges in power grids.



Effects of space weather ⁵



- many things can be affected by the space environment
- and often in many ways...




Power transformer and the Sun

before

Severe internal damage caused by the space storm of 13 March 1989

after



but be careful:
this was the only extreme case we know of...




What is space weather?




Current Space Weather

Learn more about [Space Weather](#)


NOAA/SEC Satellite Environment




GOES X-Ray Flux



SOHO/SEM EUV/X-ray Flux





Dst Geomagnetic Index Estimate






Low: $Dst > -20$ nT
Medium: -20 nT $> Dst > -50$ nT
High: -50 nT $> Dst > -100$ nT
Extreme: $Dst < -100$ nT


Auroral Activity Extrapolated from NOAA POES



SOHO/ERNE High Energy Proton Flux SOHO CELIAS/MTOF Proton Monitor ACE Solar Wind Real-Time Data



SOHO Real-time View of the Sun




<http://sohowww.nascom.nasa.gov/spaceweather/>

What questions to ask ?

8

selected physical problems to address:


the driver:



- solar irradiance
- solar (coronal) eruptions
- particle acceleration

- small & large scale structures: sunspots / faculae
- magnetic instabilities for CMEs and flares
- relativistic description of acceleration process

propagation:



- magnetic disturbances in interplanetary space
- energetic particles

- wave-particle interaction
- kinetic description of transport phenomena
- interaction of large scale solar wind/CME structures

effects on Earth:



- geomagnetic storms
- energy input into atmosphere
- energetic radiation and life
- advanced technology

- interaction of solar wind with Earth's magnetosphere
- intrusion of particles into Earth's magnetosphere
- reconnection and acceleration in magnetosphere

What questions to ask ?

selected physical problems to address:

the driver:

- solar irradiance
- solar (coronal) eruptions
- particle acceleration



alternative definition of space weather:


integration of many problems from the Sun to the Earth into an **engineering model** to predict effects on Earth.

COMPLICATION:

we have not yet understood most of the relevant individual problems...

- energetic radiation and life
- advanced technology

propag



effects







- small & large scale structures: sunspots / faculae
- magnetic instabilities for CMEs and flares
- relativistic description of acceleration process

- wave-particle interaction
- kinetic description of transport phenomena
- interaction of large scale solar wind/CME structures

- interaction of solar wind with Earth's magnetosphere
- intrusion of particles into Earth's magnetosphere
- reconnection and acceleration in magnetosphere




Solar-terrestrial relations I: solar irradiance

Fröhlich et al. (2000) SSR 94, 14
with PMOD composite d40_60_0412




Solar-terrestrial relations I: solar irradiance

Fröhlich et al. (2000) SSR 94, 14
with PMOD composite d40_60_0412



Comparing solar irradiance to Earth's temperature



Solanki et al. (2002)

Solar-terrestrial relations I: solar irradiance



total solar irradiance: $\sim 1366 \pm 1 \text{ W/m}^2$

What is the physical basis of solar variability?


How do sunspots form and evolve?

What causes “faculae” to be bright?


How do small and large structures interact?



Sunspot




faculae



Solar-terrestrial relations II: X-ray & EUV


EUV affects the Earth:

- thermosphere (heating)
- chemical reactions (e.g. ozone)
- ionization / ionosphere expansion



How does coronal heating work?


And how does it control the X-ray and EUV brightness?



1993

100 x brighter!


1991
maximum



Yohkoh Soft X-ray Telescope (SXT)

Solar–terrestrial relations III: mass ejections

- Large magnetic structures become unstable
 - coronal mass ejections (CME)
 - flares → high-energy particles
- interaction with Earth's magnetosphere
- danger to instrumentation and life (in space)



**What is the basic
physical mechanism
driving the corona unstable ?**



Northern light at the Schauinsland
(Sternfreunde Breisgau) 29.10.2003

observations:
EIT / Lasco /SOHO


sketch:
Earth's magnetosphere

Prominences and magnetic field

prominences are found above magnetic neutral lines

H α image [close up]


Magnetogram [close up]



What is a prominence ?


the „hammock“
of Kippenhahn & Schlüter (1957):

- cool dense plasma ($\sim 10^4$ K)
in a hot surrounding ($\sim 10^6$ K)
- enough (cool) plasma for significant
absorption of photospheric emission




modern idea:
complex
helical structure

over-arching
magnetic field
holding prominence down





the „hammock“
of Kippenhahn & Schlüter (1957):

- **cool dense plasma** ($\sim 10^4$ K)
in a hot surrounding ($\sim 10^6$ K)
- enough (cool) plasma for significant
absorption of photospheric emission




What is a coronal loop ?

emission of **hot plasma** ($\sim 10^6$ K)
with enhanced density as
compared to the background corona
along the magnetic field





Filaments and prominences




filament: cool plasma held by magnetic field
absorbs photospheric light

prominence: cool plasma seen in emission





1. first one prominence behind the limb seen in emission erupts
- 2 then the prominence in the front seen in absorption takes of...



almost always:

- single magnetic flux rope
- topology preserved

mostly:

- helical shape
- signature of twist

often:

- ejection (CME)
- high speeds ($\sim v_A$)

Coronal mass ejections




eruption on 4 Jan 2002

Lasco C2

rapid acceleration

Lasco C3


and huge expansion



Eruption of prominence
(seen dark in absorption)
and subsequent
brightening of
"reconnected" post flare loops




EIT 195Å / Fe XII ~1.5 MK

CME properties



huge expansion $> 10^3$
 huge solid angle $> \pi/2$
 often twisted flux ropes

two CME classes: fast & slow




High Altitude Observatory / Solar Maximum Mission


Three-part structure:

- core – prominence
- cavity – expanding flux rope?
- front – swept up plasma



A very simplified scenario for a CME




A very simplified scenario for a CME



tether cutting: “runaway” reconnection




magnetic breakout: unstable arcade triggered (& driven ?) by reconnection




Antiochos et al. (1999...)

flux cancellation at neutral line forms flux rope




Amari et al (2003)


flux rope instability



Török & Kliem (2005)




to understand the interaction with the Earth:
first understand the origin of the magnetic cloud, namely the CME ejection



Helical
kink
($m=1$)




Gerrard et al. (2001)
A&A 373, 1089









Török et al. (2004) A&A 413, L27

- kink is an ideal MHD instability → twisting a flexible tube if twist is above threshold:
 - twist "transformed" into **writhe**
- conserved: helicity \sim twist + writhe
- twist threshold: $\Phi = 2\pi N$ with $N \approx 1 \dots 2$

- many erupting filaments / prominences:
 - suggest twisted field
 - develop helical shape
- Sakurai (1976) suggested kink instability as driver of prominence eruptions
- *recent years:*
kink instability as explanation only for confined events
- *very recently:*
kink instability triggers also ejective events (CMEs)
(Török & Kliem 2005, Fan 2005)



A confined filament eruption



- one possible driver is rotational motion of foot points
 - ➔ energy stored in twist of magnetic field

- helical kink instability triggers event

HERE:


- filament eruption is confined
 - ➔ no outbreak / CME

investigate models with different

- flux rope twist
- overlying field


- ➔ strong overlying magnetic field can prevent eruption

X-ray sigmoids and flux eruption




ideal MHD simulation of flux rope eruption through kink instability

magnetic field lines



inner flux rope

current sheets



Yohkoh soft X-rays

assumption:
heating concentrated in current sheets



→ current sheets outline
expected hot X-ray emission

→ very good qualitative match
to observations

Ejective filament eruption

TRACE 1600 Å – cool ejected material
chromospheric temperatures + C IV (10^5 K)

flux rope kink instability
small overlying B allows ejection



(some) open questions for kink instability-based models


- huge expansions of CMEs by $> 10^3$
- eruptions with little or no apparent helical shape



Torus instability ?

Finally: a complete "space weather" model


Virtual space weather from the sun to Earth



Center for
Space Environment Modeling
Univ. Michigan
Gombosi, Zeeuw, Groth, Powell,
Stout, Roussev, Manchester, Toth...

The perfect storm packs the energy of 100 million atomic bombs

Solution at 9 hours




April, 1999

Simulating space weather: numerical challenge

needed for these global models:

advanced codes for many different physical problems:

- adaptive mesh refinement (AMR) to resolve large and small scales
- MHD codes
- particle codes
-



Physics Domain

Models / Codes

Solar Corona

BATSRUS

Eruptive Event Generator

BATSRUS

Inner Heliosphere

BATSRUS

Solar Energetic Particles

Kóta's SEP model

Global Magnetosphere

BATSRUS

Inner Magnetosphere

Rice Convection Model

Ionosphere Electrodynamics

Ridley's potential solver


Upper Atmosphere

General Ionosphere-Thermosphere Model (GITM)


Bats'r'us:
snapshot of
grid (AMR)
for CME model

modules of the space weather code of
Center for Space Environment Modeling,
University of Michigan


CME eruption and propagation to Earth



Interaction with Earth



just before the CME
hits the Earth



Earth's magnetosphere
after the interaction

The global space weather model puts together many modules:
good "engineering model" of physical phenomena

- a general problem: **not yet in real time**
 - many weeks to simulate an event which last for only some days...
- there are important physics pieces still missing!
for the coronal parts:
 - solar wind heating and acceleration
 - problem of CME initiation
 - reconnection processes
- spatial resolution in corona:
 - currently AMR with smallest cells $1/32 R_{\odot} = 21 \text{ Mm}$ (!)
 - this resolution certainly cannot catch the relevant physics
 - for comparison: coronal box models: computational domain $\sim 60 \times 60 \times 40 \text{ Mm}$

However: if one is interested in an engineering approach
i.e. only predict when, where and how a CME hits the Earth
this might be an appropriate approach

- there are many ways in which the Sun affects the Earth
 - Luminosity: bolometric, X-rays, VUV etc.
 - particle radiation: CMEs, energetic particles
 - magnetic field: cosmic rays
- the most relevant phenomenon concerning corona: CME
 - different scenarios for CME initiation
 - instabilities, tether cutting, breakout...
 - all scenarios are (in the end) driven by photospheric shuffling of magnetic field
- global models of CME initiation to Earth interaction needed for "space weather"
 - global models currently in an "engineering state"
 - detailed physics CME and/or interaction with Earth are not really included