

An artistic rendering of a Cluster mission satellite in space. The satellite is cylindrical with a purple base and a gold-colored top section. It has several long, thin antennas extending from it. In the background, the Earth is visible on the left, and a bright, glowing sun is on the right. Other satellites are visible in the distance, connected by thin lines. The text 'The Cluster Mission to the Earth's Magnetosphere' is written in yellow, italicized font across the top. The name 'Patrick W. Daly' is written in white below the title. The 'MPS' logo is in the bottom center.

The Cluster Mission to the Earth's Magnetosphere

Patrick W. Daly

MPS

- 1 The Mission
 - Aims
 - Concept
 - History
 - Current Status
- 2 Payload
- 3 The RAPID Instrument
 - Electrons
 - Ions
 - 3-D plots
 - Whole Magnetosphere
 - Data Transmission
 - Data Processing

The Mission

Aims
Concept
History
Status

Payload

RAPID

Electrons
Ions
3-D plots
Whole MagSphere
Data Transmission
Data Processing

The Cluster Mission

The Cluster mission is an in-situ investigation of the Earth's magnetosphere using **four identical spacecraft** simultaneously.

Advantages

- accurate determination of three-dimensional and time-varying phenomena
- distinguishing between spatial and temporal variations

The Mission

Aims
Concept
History
Status

Payload

RAPID

Electrons
Ions
3-D plots
Whole MagSphere
Data Transmission
Data Processing

Cluster's main goal is to study the small-scale plasma structures in space and time in the key plasma regions:

- solar wind and bow shock
- magnetopause
- polar cusp
- magnetotail
- auroral zone

The Mission

Aims

Concept
History
Status

Payload


RAPID

Electrons
Ions
3-D plots
Whole MagSphere
Data Transmission
Data Processing

Example: Cusp

The interaction between the **solar wind** and the **magnetosphere** is a key element in the ISTP program.

Example: direct entry of solar wind particles through the polar cusps, which act as two magnetic funnels, one in each hemisphere, focusing the solar wind particles (rather like a telescope) on photons.



The Mission

Aims

- Concept
- History
- Status

Payload

RAPID

- Electrons
- Ions
- 3-D plots
- Whole MagSphere
- Data Transmission
- Data Processing

Example: Substorms

Another example: acceleration of plasma in the magnetotail during substorms.

A reversal of the interplanetary magnetic field from north to south can initiate such a substorm, releasing a large quantity of particles towards the Earth.

Both entry mechanisms produce auroras when the precipitating electrons and ions are absorbed in the neutral atmosphere.

These particles can have dramatic effects on, e.g.:

- power transmission and telecommunications,
- satellite operations in geostationary orbit.

Predicting such *space weather* effects is also one of the goals of the Cluster Mission.

The Mission

Aims

Concept

History

Status

Payload

RAPID

Electrons

Ions

3-D plots

Whole MagSphere

Data Transmission

Data Processing

The Mission

Aims

Concept

History

Status

Payload

RAPID

Electrons

Ions

3-D plots

Whole MagSphere

Data Transmission

Data Processing

The fundamental idea behind Cluster:

- Four **identical** satellites in a tetrahedron formation
- with identical payloads designed for magnetospheric plasma studies:
 - electromagnetic waves
 - electric and magnetic fields
 - electron and ion measurements over broad energy spectrum
- on elliptical polar orbit to cover all the regions magnetosphere in course of one year (4-19 R_E)

Cluster has a long story behind it, going back to the early 1980's.

- Cluster was proposed as an official ESA Mission in **1984**, and model payloads and technical/financial feasibilities were worked out.
- Announcement of Opportunity for experimenters came out in **1987**.
- Proposals accepted, begin building experiments in **1989**.
- Integration, tests in **1994-1996**.
- Launch with first Ariane 5 rocket on **June 4, 1996**.
- End of Cluster (I) Mission, **June 4, 1996**.

The Mission

Aims
Concept
History
Status

Payload

RAPID

Electrons
Ions
3-D plots
Whole MagSphere
Data Transmission
Data Processing

In spite of great financial and technical difficulties, ESA and the participating institutes decided that Cluster should be rebuilt.

- Final decision to go ahead in **November 1996**.
- Spacecraft and experiments rebuilt **exactly as before**, as far as that was possible.
- New start for Cluster II on two Soyuz rockets, **July, August 2000**.
- This time, **successful**.



The Mission

Aims

Concept

History

Status

Payload

RAPID

Electrons

Ions

3-D plots

Whole MagSphere

Data Transmission

Data Processing

- After one extension to the end of 2005, the mission has now been extended to the end of 2009. (Reentry will occur during 2010.)
- Most of the experiments are still working satisfactorily, although some degradation and failures have occurred.
- The spacecraft are still functioning well, fuel is sufficient, solar cells acceptable, but the batteries will likely not last much longer. (This means reduced operations during eclipses.)
- Spacecraft separations have been varied from 200 to 5000 km.
- As of July 2005, permanent large separation of 10 000 km between 3 spacecraft; 4th one can vary 100 to 10 000 km. This is called multi-scaling.

The Mission

Aims

Concept

History

Status

Payload

RAPID

Electrons

Ions

3-D plots

Whole MagSphere

Data Transmission

Data Processing

The Payload

The Cluster satellites each contain 11 different experiments.

Wave Instruments

DWP coordinates the other wave instruments and performs particle/wave correlations.

EFW measures the electric field in the plane \perp spin axis and its power density up to 180 Hz.

STAFF measures magnetic field fluctuations $<$ 4 kHz.

WHISPER measures electron density with a relaxation sounder, and electric fields in frequency range 2–80 kHz.

WBD provides electric or magnetic waveforms up to 577 kHz.

The Mission

Aims

Concept

History

Status

Payload

RAPID

Electrons

Ions

3-D plots

Whole MagSphere

Data Transmission

Data Processing

Fields

ASPOC controls the spacecraft potential by emitting indium ions, preventing charging which would hamper the electron instruments.

FGM fluxgate magnetometer, measures the magnetic field in 3 dimensions.

EDI measures electric fields by shooting keV electrons into space and detecting their return to the spacecraft via gyromotion.

The Mission

Aims
Concept
History
Status

Payload

RAPID

Electrons
Ions
3-D plots
Whole MagSphere
Data Transmission
Data Processing

Particles

CIS hemispherical electrostatic ion spectrometer, in 4π directions, up to 32 keV/e, for H^+ , He^+ , He^{++} , and O^+ .

PEACE two hemispherical electrostatic electron spectrometers, for 4π coverage up to ~ 26 keV.

RAPID measures energetic electrons and ions with imaging techniques and time-of-flight methods, from ~ 30 –400 keV, with 4π coverage.

Payload: The Movie

Cluster Mission

P. W. Daly


The Mission

Aims
Concept
History
Status

Payload

RAPID

Electrons
Ions
3-D plots
Whole MagSphere
Data Transmission
Data Processing



The Mission

Aims
Concept
History
Status

Payload

RAPID

Electrons
Ions
3-D plots
Whole MagSphere
Data Transmission
Data Processing

RAPID stands for

*Research with Adaptive Particle Imaging
Detectors*

and is an energetic ion and electron ($E > 30$ keV) imaging spectrometer.

The IES electron heads

The Mission


Aims
Concept
History
Status

Payload

RAPID

Electrons

Ions
3-D plots
Whole MagSphere
Data Transmission
Data Processing



Each of the electron heads consists of

- a “pin-hole” aperture
- an absorber for ions ($E < 350$ keV)
- 3 solid state detectors.

The 3 detectors determine the electron incidence angle to within 20° .

The IIMS/SCENIC ion heads


The Mission

Aims
Concept
History
Status

Payload


RAPID

Electrons
Ions
3-D plots
Whole MagSphere
Data Transmission
Data Processing



Particles entering the ion heads must first pass through the collimating plates before encountering the time-of-flight arrangement.

The Time-of-Flight Measurement



One of 3 IIMS (Ions) Sensors

The collimated ions encounter a thin foil, emitting electrons that serve as the **START** signal.

The **STOP** signal is the absorption in the solid state detector that determines the ion's energy.

The **START** signal also serves to localize the incident direction within 15°.

The combination of energy (SSD) and velocity (TOF) determine the ion mass and thus identify its species.

The Mission

Aims
Concept
History
Status

Payload

RAPID

Electrons

Ions

3-D plots

Whole MagSphere

Data Transmission

Data Processing

RAPID Specifications

The Mission

Aims
Concept
History
Status

Payload

RAPID

Electrons

Ions

3-D plots

Whole MagSphere

Data Transmission

Data Processing

Energy ranges:	Hydrogen	28–1500 keV
	Helium	29–1500
	CNO	92–1500
	Electrons	26–400
Mass range:		1, 4, 12–16, 28–56
Resolution (A/dA):	Oxygen	4
Field-of-view:	IIMS (Ions)	$\pm 3^\circ \times 180^\circ$
	IES (Elec.)	$\pm 17.5^\circ \times 180^\circ$
Geometry factor: (for 180°)	IIMS	$2.6 \times 10^{-2} \text{ cm}^2 \cdot \text{sr}$
	IES	2.0×10^{-2}

Angular Coverage in 3-D

The Mission

Aims
Concept
History
Status


Payload

RAPID

Electrons
Ions

3-D plots

Whole MagSphere
Data Transmission
Data Processing



Note: the spin axis is directed towards the *southern* ecliptic pole!

The 3-D plots

The Mission

Aims
Concept
History
Status


Payload

RAPID

Electrons
Ions

3-D plots

Whole MagSphere
Data Transmission
Data Processing



RAPID protons (30 keV)

Possible formats:

- 1 **Spacecraft**
- 2 **GSE coordinates**
- 3 **Bi-spherical GSE**
- 4 **Smoothed**

The Mission

- Aims
- Concept
- History
- Status


Payload

RAPID

- Electrons
- Ions

3-D plots

- Whole MagSphere
- Data Transmission
- Data Processing



The plots on the following slides were produced by

Jackie Davies
Rutherford Appleton Laboratory

They show RAPID electron and proton flux intensities from June 2002 to January 2004, against location in the magnetosphere.

(These are so-called *Bryant* plots.)

Bryant Plot

The Mission

- Aims
- Concept
- History
- Status


Payload

RAPID

- Electrons
- Ions
- 3-D plots

Whole MagSphere

- Data Transmission
- Data Processing



MSP V12.0.2, reference spacecraft = 1

Bryant plot software V4.2, Tue Jun 28 14:02:37 2005

RAPID Electrons

Cluster Mission

P. W. Daly

The Mission

Aims
Concept
History
Status


Payload

RAPID

Electrons
Ions
3-D plots

Whole MagSphere


Data Transmission
Data Processing



RAPID Protons

Cluster Mission

P. W. Daly



The Mission

- Aims
- Concept
- History
- Status

Payload

RAPID

- Electrons
- Ions
- 3-D plots

Whole MagSphere

- Data Transmission
- Data Processing

Cluster Magnetic Azimuth

Cluster Mission

P. W. Daly

The Mission

Aims
Concept
History
Status


Payload

RAPID

Electrons
Ions
3-D plots

Whole MagSphere

Data Transmission
Data Processing



On board

- Once per spin (~ 4 s), the RAPID DPU stores the accumulated counts into an **Experiment Data Block**, EDB. This contains 512 bytes (or 2304 bytes in high rate mode).
- The counts are compressed: one byte encodes numbers from 0 to 734000.
- Once per spacecraft cycle of 5.1522 s, a block of 660 bytes (2976 in high rate) is shifted to the output stream.
- The data are stored on board until contact with the receiving stations, when they are downloaded.
- Accurate planning is needed to avoid storage overflow and data loss.

The Mission

Aims
Concept
History
Status

Payload

RAPID

Electrons
Ions
3-D plots
Whole MagSphere
Data Transmission
Data Processing

On the Ground

- Data received by ground station.
- Transmitted to ESOC in Darmstadt.
- Reconstructed time tags added.
- Data are put online for Experiment Teams to retrieve.

Data Disposition

- Data available 20 days at ESOC.
- PI Teams order the data that they wish.
- Ordered data transferred per FTP to the recipient.
- Data are recorded to CDs on daily basis (2-3 CDs per day).
- Sets of CDs are mailed to PIs, Data Centres, Co-institutes.

The Mission

Aims
Concept
History
Status

Payload

RAPID

Electrons
Ions
3-D plots
Whole MagSphere
Data Transmission
Data Processing

Raw Data

- Raw data are repackaged, merging filtered science and HK data to form **Merged Science Files**, MSF.
- The MSF files, available to Co-Is and CAA, are primary input for further processing.

Science Data

- Science data generated with a program `msf2sci`.
- It reads the raw data, locates the EDBs within the transmission packages, reconstructs the original accumulation time stamp, . . .
- applies calibration parameters, determines energy levels, accumulation time intervals, etc, . . .
- and writes flux values to SCI file in an ascii format.
- The SCI files are inputs to further plotting and analysis programs.

The Mission

Aims
Concept
History
Status

Payload

RAPID

Electrons
Ions
3-D plots
Whole MagSphere
Data Transmission
Data Processing