

Planetary Interiors

Ulrich Christensen



Earth as a prototype planet

Informations from shape, gravity and rotation

Internal structure of terrestrial planets and icy moons

The interior of gas planets

Prototype planet Earth



Most knowledge on Earth's internal structure comes from seismology (body waves and free oscillations). It provides information on elastic properties (incompressibility k , shear modulus μ) and density ρ as function of radius. Basic parts of the Earth are the **crust** (6-50 km thick), the solid **mantle** (to 2900 km depth), the liquid outer **core**, and a small solid inner core.

We assume that the internal structure of other terrestrial planets is basically similar,

Composition of different parts of the Earth




Sources of information:

Crust – plenty of direct samples
 Upper Mantle – samples from exposed mantle rock or xenoliths (solid mantle rock carried upwards in volcanic vents)
 Deep mantle and core – indirect

	Continental crust(0.2%)	Oceanic crust(0.1%)	Mantle (68%)	Core (32%)
SiO ₂	60%	50%	46%	
MgO	3 %	8 %	38%	
FeO	4%	9%	8%	Fe: 85 %
Al ₂ O ₃	17%	16%	4%	
CaO	7%	12%	3%	
Na ₂ O	3%	1%	<1%	
Rock type:	Granite	Basalt	Peridotite	Iron alloy
Minerals:	Quartz SiO ₂ Feldspar: CaAl ₂ Si ₂ O ₈ – NaAlSi ₃ O ₈ (Plagioclase)	Plagioclase Pyroxene: CaAlSi ₂ O ₆ – (Mg,Fe)SiO ₃	Olivine: (Mg,Fe) ₂ SiO ₄ Pyroxene, Garnet: Mg ₃ Al ₂ Si ₃ O ₁₂	

Mantle rock is rich in magnesium (over 90% consists of Si, Mg, Fe, O). Crustal rocks contain little Mg and comparatively more Ca and Al. The core must be made mostly of iron, because Fe is the only element that (1) has the right density and compressibility at high pressure to satisfy the seismological data and (2) is sufficiently abundant. The core density is slightly less dense than iron at core pressures ⇒ 5-10% of light elements (Si, S, O) must be present.

Information on internal structure from shape, gravity field and rotation



Seismological information is available only for the Earth and in limited amounts for the Moon. Various geodetic data put constraints on the internal structure, but the ambiguity is much larger than for seismic data.

Gravity field: fundamentals

Gravitational potential V , Gravity (acceleration) $\mathbf{g} = -\text{grad } V$

Point mass: $V = -GM/r$ (also for spherically symmetric body)

Ellipsoid: $V = -\frac{GM}{r} \left(1 - J_2 \left(\frac{a}{r}\right)^2 P_2(\cos \theta) + \dots \right)$ $P_2(\cos \theta) = \frac{3}{2} \cos^2 \theta - \frac{1}{2}$

General: $V = -\frac{GM}{r} \left(1 + \sum_{\ell=2}^{\infty} \left(\frac{a}{r}\right)^{\ell} \sum_{m=0}^{\ell} P_{\ell}^m(\cos \theta) [C_{\ell}^m \cos m\varphi + S_{\ell}^m \sin m\varphi] \right)$

General description of gravity field in terms of spherical harmonic functions. Degree $\ell=0$ is the monopole term, $\ell=2$ the quadrupole, $\ell=3$ the octupole, etc. A dipole term does not exist when the coordinated system is fixed to the centre of mass. J , C , S are non-dimensional numbers. Note: $J_2 = -C_2^0$ (times a constant depending on the normalization of the P_{ℓ}^m)

Symbols [bold symbols stand for vectors]: G – gravitational constant, M – total mass of a body, r – radial distance from centre of mass, a – reference radius of planet, e.g. mean equatorial radius, θ – colatitude, φ – longitude, P_n – Legendre polynomial of degree n , P_{ℓ}^m – associated Legendre function of degree ℓ and order m , J_n , C_{ℓ}^m , S_{ℓ}^m – expansion coefficients

Measuring the gravity field


- Without a visiting spacecraft, the monopole gravity term (the mass M) can be determined by the orbital perturbations on other planetary bodies or from the orbital parameters of moons (if they exist)
- From a spacecraft flyby, M can be determined with great accuracy. J_2 and possibly other low-degree gravity coefficient are obtained with less accuracy
- With an orbiting spacecraft, the gravity field can be determined up to high degree (Mars up to $\ell \approx 60$, Earth up to $\ell \approx 180$)
- The acceleration of a spacecraft orbiting (or passing) another planet is determined with high accuracy by radio-doppler-tracking: The Doppler shift of the carrier frequency used for telecommunication is proportional to the line-of-sight velocity of the spacecraft relative to the receiving antenna. Δv can be measured to much better than a mm/sec.
- On Earth, direct measurements of g at many locations complement other techniques.
- The ocean surface on Earth is nearly an equipotential surface of the gravity potential. Its precise determination by laser altimetry from an orbiting S/C reveals small-wavelength structures in the gravity field.

Mean density and uncompressed density

From the shape (volume) and mass of a planet, the mean density ρ_{mean} is obtained. It depends on chemical composition, but by self-compression also on the size of the planet (and its internal temperature, though in case of terrestrial planets only weakly). In order to compare planets of different size in terms of possible differences in composition, an **uncompressed density** can be calculated - the mean density it would have, when its material were at 1 bar. This requires knowledge of incompressibility k (from high-pressure experiments or from seismology in case of the Earth).

	ρ_{mean}	$\rho_{\text{uncompressed}}$	
Earth	5515	4060	kg m^{-3}
Moon	3341	3315	kg m^{-3}

The mean density alone gives no clue on the radial distribution of density: a body could be an undifferentiated mixture (e.g. of metal and silicate, or of ice and rock in the outer solar system), or could have separated in different layers (e.g. mantle and core).



Moment of inertia


$L = I\omega$ L : Angular momentum, ω : angular frequency, I moment of inertia

$I = \iiint \rho s^2 dV$ for rotation around an arbitrary axis, s is distance from that axis

I is a symmetric tensor. It has 3 principal axes and 3 principal components (maximum, intermediate, minimum moment of inertia: $C \geq B \geq A$.) For a spherically symmetric body rotating around polar axis

$C = \frac{8\pi}{3} \int_0^a \rho(r) r^4 dr$ compare with integral for mass $M = 4\pi \int_0^a \rho(r) r^2 dr$

In planetary science, the maximum moment of a nearly radially symmetric body is usually expressed as $C/(Ma^2)$, a dimensionless number. Its value provides information on how strongly the mass is concentrated towards the centre.



Symbols: L – angular momentum, I moment of inertia (C, B, A – principal components), ω rotation frequency, s – distance from rotation axis, dV – volume element, M – total mass, a – planetary radius (reference value), a_c – core radius, ρ_m – mantle density, ρ_c – core density

Determining planetary moments of inertia

McCullagh's formula $J_2 = \frac{C - \frac{1}{2}(B + A)}{Ma^2}$ for ellipsoid (B=A): $J_2 = \frac{C - A}{Ma^2}$

In order to obtain $C/(Ma^2)$, the dynamical ellipticity $H = (C-A)/C$ is needed. It can be uniquely determined from observation of the precession of the planetary rotation axis due to the solar torque (plus lunar torque in case of Earth) on the equatorial bulge. For solar torque alone, the precession frequency relates to H by:


$$\omega_p = \frac{3}{2} \frac{\omega_{\text{orbit}}^2}{\omega_{\text{spin}}} H \cos \varepsilon$$

When the body is in a locked rotational state (Moon), H can be deduced from nutation.

For the Earth $T_p = 2\pi/\omega_p = 25,800$ yr (but here also the lunar torque must be accounted)

$$H = 1/306 \text{ and } J_2 = 1.08 \times 10^{-3} \Rightarrow C/(Ma^2) = J_2/H = 0.3308.$$


This value is used, together with free oscillation data, to constrain the radial density distribution.



Symbols: J_2 – gravity moment, ω_p precession frequency, ω_{orbit} – orbital frequency (motion around sun), ω_{spin} – spin frequency, ε - obliquity

Determining planetary moments of inertia II

For many bodies no precession data are available. If the body rotates sufficiently rapidly and if its shape can be assumed to be in hydrostatic equilibrium [i.e. equipotential surfaces are also surfaces of constant density], it is possible to derive $C/(Ma^2)$ from the degree of ellipsoidal flattening or the effect of this flattening on the gravity field (its J_2 -term). At the same spin rate, a body will flatten less when its mass is concentrated towards the centre.



Darwin-Radau theory for an slightly flattened ellipsoid in hydrostatic equilibrium

$$m = \frac{\omega_{\text{spin}}^2 a^3}{GM} \quad \text{measures rotational effects (ratio of centrifugal to gravity force at equator).}$$



Flattening is $f = (a-c)/a$. The following relations hold approximately:

$$f = \frac{3}{2} J_2 + \frac{1}{2} m \qquad \frac{C}{Ma^2} = \frac{2}{3} - \frac{4}{15} \sqrt{\frac{5m}{2f} - 1} \qquad \frac{C}{Ma^2} = \frac{2}{3} - \frac{4}{15} \sqrt{\frac{4m - 3J_2}{m + 3J_2}}$$

Symbols: a – equator radius, c- polar radius, f – flattening, m – centrifugal factor (non-dimensional number)

Structural models for terrestrial planets

	ρ_{mean} kg m ⁻³	ρ_{uncompr} kg m ⁻³	C/Ma ²
Mercury	5430	5280	?
Venus	5245	3990	?
Earth	5515	4060	0.3308
Moon	3341	3315	0.390
Mars	3935	3730	0.366




Assuming that the terrestrial planets are made up of the same basic components as Earth (silicates / iron alloy with zero-pressure densities of 3300 kg m⁻³ and 7000 kg m⁻³, respectively), core sizes can be derived.

Ambiguities remain, even when ρ_{mean} and C/Ma² are known: the three density models for Mars satisfy both data, but have different core radii and densities with different sulphur contents in the core.

Interior of Galilean satellites

	ρ [kg m ⁻³]	C/Ma ²
Io Silicate mantle Iron core	3530	0.378
Europa (Thin) ice layer Silicate mantle Iron core	3020	0.347
Ganymede Thick ice shell Silicate mantle Iron core	1940	0.311
Callisto Ice layer Ice/silicate/iron mixture below	1850	0.358



From close Galileo flybys
mean density and J_2 (assume
hydrostatic shape $\Rightarrow C/(Ma^2)$)

Low density of outer satellites
 \Rightarrow substantial ice (H_2O)
component.

Three-layer models (ice, rock,
iron) except for Io. Assume
rock/Fe ratio.

Callisto's C/Ma^2 too large for
complete differentiation \Rightarrow
core is probably an undifferen-
tiated rock-ice mixture.

Gravity anomalies

Gravity anomaly Δg : Deviation of gravity (at a reference surface) from theoretical gravity of a rotating ellipsoidal body.


Geoid anomaly ΔN : Deviation of an equipotential surface from a reference ellipsoid (is zero for a body in perfect hydrostatic equilibrium).

Isostasy means that the extra mass of topographic elevations is compensated, at not too great depth, by a mass deficit (and vice versa for depressions).





In the Airy model, a crust with constant density ρ_c is assumed, floating like an iceberg on the mantle with higher density ρ_m . A mountain chain has a deep crustal root.

When the horizontal scale of the topography is much larger than the vertical scales, the gravity anomaly for isostatically compensated topography is approximately zero.



Without compensation, or for imperfect compensation, a gravity anomaly is observed. Also for very deep compensation (depth not negligible compared to horizontal scale) a gravity anomaly is found.



Examples for different tectonic settings



Gravity field of Mars



Comparing the gravity anomaly of Mars with the topographic height, we can conclude:

Tharsis volcanoes not compensated, only slight indication for elastic plate bending
⇒ volcano imposed superficially on thick lithosphere

Hellas basin shows small gravity anomaly
⇒ compensated by thinned crust



Valles Marineris not compensated

Topographic dichotomy (Southern highland, Northern lowlands) compensated
⇒ crustal thickness variation


Tharsis bulge shows large-scale positive anomaly. Because incomplete compensation is unlikely for such broad area, deep compensation must be assumed (for example by a huge mantle plume)

Properties of Jupiter and Saturn

	Jupiter	Saturn
Equatorial radius [km]	71,500	60,300
Flattening	1 / 15.7	1 / 9.8
Mass / Earth mass	318	95
Mean density [kg m⁻³]	1330	690
Rotation period [h]	9.9	10.7
Equatorial gravity [ms⁻²]	22.9	9.1
Surface temperature [K]	124	95
Emitted/absorbed power	1.7	1.8
Atmospheric composition		
H₂	0.85	0.94
He	0.15	0.05?
H₂O	0.001	0.001
CH₄	0.001	0.002
NH₃	0.002	0.001




Transition to metallic hydrogen



Quantum mechanical calculations suggested that solid or fluid hydrogen becomes metallic at high pressure, i.e. has free electrons and becomes a good electrical conductor.



In shock wave experiments it was possible to measure the conductivity. The experiments show that metallization occurs between 1 and 1.4 Mbar (100 – 140 GPa).

Phase diagram of hydrogen



The temperature and pressure conditions in the gas planets are included, calculated with the assumption of an adiabatic temperature gradient starting at the observed surface temp.

Interior of Jupiter




Interior models for Jupiter (and Saturn) are based on the phase diagram and a theoretical equation of state - $\rho(p,T)$ - for hydrogen-helium mixtures and the assumption of adiabatic temperature variation. They are constrained by the requirement to fit the observed flattening and the gravity coefficients J_2 , J_4 .

A small rocky core of a few Earth masses is not required by the geophysical data for Jupiter (but for Saturn), but is likely in any case, because a sizeable solid nucleus may be required to be present first in order to accrete H_2 and He onto it by gravitation.

Interior of Saturn

SATURN MODELS



The principal structure of Saturn is similar to that of Jupiter. Because pressure is lower, the boundary between metallic and molecular hydrogen is deeper (half the planetary radius) and the rocky core may be bigger (14 Earth masses).

An important difference may be that at the top of the metallic hydrogen region, He becomes immiscible in Saturn, where temperatures are lower than in Jupiter (compare slide 5.4).

In this region, He may separate and form rain drops that sink and dissolve again at greater depth in the metallic H-region.


- ⇒ He becomes depleted also in the molecular hydrogen region, which mixes with the top of the metallic region
- ⇒ Extra gravitational energy is released. This could explain the large excess luminosity of Saturn, which is more than what theoretical models based on cooling and contraction predict.

Interior of Uranus and Neptune

Uranus and Neptune cannot be made mostly of H + He which dominate in the atmosphere. Because of their low mass (lower gravitation), in that case the mean density would be much lower than that of Saturn.

The region below 70% of the planetary radius is a supercritical fluid consisting mainly of H₂O (plus NH₃ and CH₄). Ionic electrical conductivity allows for a dynamo to operate in this region and generate the observed magnetic fields.

A rocky core has probably around one Earth mass.



Mainly H₂+He

„Ices“ (H₂O, NH₃, CH₄)

Rocky core (~ 1 M_{earth})